


Nutrition for PLWHA - Counselling Cards



Nutrition for PLWHA - Counselling Cards is a publication of Regional Centre for Quality of Health Care (RCQHC), Kampala - Uganda. Technical and financial support for development of this publication was provided by the Food and Nutrition Technical Assistance (FANTA) and LINKAGES projects.

RCQHC is a regional quality of health care capacity development institute largely supported by REDSO/ESA in Nairobi and Makerere University, Kampala.

The FANTA Project is supported by the Regional Economic Development Service Office/East and Southern Africa (REDSO/ESA) and the Office of Health, Infectious Disease and Nutrition of the Bureau for Global Health at the U.S. Agency for International Development (USAID), under terms of Cooperative Agreement No. HRN-A-00-98-00046-00 awarded to the Academy for Educational Development (AED).

The LINKAGES Project is supported by the Regional Economic Development Service Office/East and Southern Africa (REDSO/ESA) and GH/HIDN of the United States Agency for International Development (USAID), under the terms of Cooperative Agreement No. HRN-A-00-97-00007-00.

The opinions expressed herein are those of the authors and do not necessarily reflect the views of USAID.

March 2005 (revised)



For copies of this flip chart contact:

Regional Centre for Quality of Health Care, Makerere University Medical School,
P O Box 29140, Kampala -Uganda, Tel. 256-41-530888, Fax 256 530876.

Acknowledgments

This publication is a product of the Regional Centre for Quality of Health Care (RCQHC). A number of people and organisations contributed to the development of this booklet. The content was drafted by the RCQHC, the FANTA Project, the LINKAGES Project, Mary Materu of CONSENUTH in Tanzania and Gakenia Wamuyu Maina of IPH. Additional comments were given during a national workshop held to review the materials in Jinja, May 2004.

Appreciation is expressed for technical input from participants of the Jinja 2004 workshop, specifically, Karen Kyampaire Apophia and Denis Okwar of ACDI/VOCA; Sande Julius of AIC Uganda; Phoebe Nabongo of Infectious Disease Institute, Mulago; Mulindwa Rwabaingi of Joint Clinical Research Centre; Mary Butamanya and Sheila Nyakwezi of Mildmay; Ursula Wangwe of Ministry of Health; Magero Jacinta of NACWOLA; Faith Akiiki of National Forum of PLWHA; Maria Nanyonga and Resty Ingabire of Nsambya Home Based Care; Joy Elizabeth Ogutu of PATCH CONSULT; Samalie Bananuka of STD/ACP Ministry of Health; Penninah Namusi, Hellen Ekideit and Juliana Nyombi of TASO; Peter Ddamulira of THETA; Lilian Nakato of Uganda HIV/AIDS Control Project; and Irene Kiiza of Livelihood Security Program- World Vision. Gratitude is also due to Dr. Joel Okullo, Director of the RCQHC in Uganda for the leadership provided in helping us produce this booklet.

The RCQHC also appreciates the technical input and review comments provided by Robert Mwadime, Tony Castleman, Lora Iannotti, Dorcas Lwanga, Sandra Remancus and Liz Thomas of AED.

Table of Contents

INTRODUCTION

Theme 1 - Good nutrition is important for the well-being of PLWHA p.2

- 1.1 Good nutrition keeps the body strong
- 1.2 Poor nutrition quickens progression of HIV to AIDS
- 1.3 Good nutrition delays progression of HIV to AIDS

Theme 2 - Eating well for PLWHA p.8

- 2.1 Eat meals that have a variety of food types
- 2.2 Eat fortified foods to increase intake of essential nutrients
- 2.3 Increase energy intake by eating more
- 2.4 Eat snacks to increase energy intake
- 2.5 Address constraints to eating well

Theme 3 - Preventing infections p.18

- 3.1 Keep surroundings clean
- 3.2 Practice food hygiene and safety
- 3.3 Have regular check-ups

Theme 4 - Maintaining physical activity p.24

- 4.1 Exercise regularly and continue daily activity

Theme 5 - Manage diet related HIV/AIDS symptoms p.26

- 5.1 Loss of appetite
- 5.2 Depression and stress
- 5.3 Mouth sores and oral thrush
- 5.4 Tuberculosis
- 5.5 Anaemia
- 5.6 Diarrhoea, nausea and vomiting

INTRODUCTION

This is a flip chart containing counselling cards. These cards are designed for counselors and health workers to enhance nutrition counselling for PLWHA.

Use these cards when counselling PLWHA or their caregivers to help them:

- Make informed choices for improving their nutrition.
- Identify and use locally available foods to meet their nutritional needs.
- Understand and adopt practices that promote a healthier nutritional status.
- Understand how to use diet to manage common HIV/AIDS related conditions.

How to use the cards

1. You can use these cards for both individual and group counselling.
2. Read all the cards to know the content before using them. This will help you to select the appropriate cards in a given counselling situation.
3. You can place the flip chart on a table or hold it to your chest. The side with the illustration should face the client/s. The side with the notes should face the counselor. Hold the flip chart in a way that allows clear viewing of illustrations by the client/s.
4. After using a card, flip it over so that the client sees a new illustration. This enables the counselor to see the notes on the back of the previous page.
 - Use theme 1 cards to provide general counselling on nutrition and HIV/AIDS for both individual and group counselling.
 - Use theme 2 cards to provide specific counselling on eating well.
 - Use theme 3 cards to provide counselling on how to avoid infections.
 - Use theme 4 cards to provide counselling for PLWHA on the importance of physical activity.
 - Use theme 5 cards to provide counselling for PLWHA with HIV related symptoms

5. Maintain eye contact when talking with the client during counselling.

- Build on what the client knows. Use the key messages in the cards to reinforce or correct the client's responses as needed.
- Review with your client the key points discussed on each card to ensure the client has understood the discussion correctly.

Remember:


- Every client is unique. How he or she responds to information may vary.
- Clients with a recurring condition requiring the same nutrition intervention may have underlying medical needs which necessitate medical approaches.
- All cards may not be necessary for each client session. Select cards to use according to the individual client's needs.

Importance of nutrition counselling for PLWHA

Malnutrition worsens the effects of HIV infection by weakening the body's ability to fight diseases. On the other hand, HIV infection increases nutritional needs and vulnerability to malnutrition. Frequent and chronic illnesses compromise food production capacity, and food intake, absorption and utilization.


Improving nutrition can help to strengthen the body's ability to fight diseases and delay progression from HIV to AIDS. This makes it possible for the PLWHA to stay productive and live longer.

Good nutrition keeps the body strong.




Nutritional needs well met
• Adequate food intake,
• Balanced diet

2



Good Nutritional Status
• Weight gain,
• Muscle size maintained,
• Strong body,

HIV/AIDS



3

Strengthened ability to fight HIV and other infections

4



Reduced vulnerability to infections
Slower progression to AIDS

THEME 1 GOOD NUTRITION IS IMPORTANT FOR THE WELL-BEING OF PLWHA

1.1 GOOD NUTRITION KEEPS THE BODY STRONG

Ask the client to describe what he/she sees in the picture.
Building on the client's response, explain how good nutrition affects HIV/AIDS.

1. The picture to the left

- HIV increases the body's needs for food.
- To meet the increased food needs, the PLWHA is eating well and absorbs foods required by the body.

2. The picture above

- Because the PLWHA is able to meet his extra food needs, he will not lose weight. He will be able to stay strong and well nourished.
- The PLWHA has a well nourished body that enables him to build strength to fight HIV/AIDS and other diseases.

3. The picture far right


- The body's defence system is strengthened against disease and infection because the body has enough nutrients stored.
- The body can therefore fight off infections better. This delays progression of HIV to AIDS.

4. The picture below


- The body does not easily get infections. This makes the PLWHA stay strong and less dependent on others.
- Since the PLWHA does not get frequent infections, progression of HIV to AIDS is delayed. The PLWHA will have a good appetite and the cycle continues.

Poor nutrition quickens progression to AIDS.

- 1**
Increased Nutrition Needs
- Due to poor absorption of nutrients
 - In order to address infection and viral replication
 - Due to symptoms



- 2**
Poor Nutrition
- Weight loss
 - Muscle wasting
 - Weakness
 - Micronutrient deficiency



- 3**
Weakened ability of the body to fight HIV and other infections

HIV/AIDS



- 4**
Increased vulnerability to infections, including TB and flu, and thus faster progression to AIDS

THEME 1

GOOD NUTRITION IS IMPORTANT FOR THE WELL-BEING OF PLWHA

1.2 POOR NUTRITION QUICKENS PROGRESSION OF HIV TO AIDS

Ask the client what he/she sees in the picture.

Explain to the client how poor nutritional status affects progression of HIV to AIDS.

1. The picture to the left

- HIV and frequent infections increase the nutritional needs of the PLWHA.
- But the PLWHA cannot take in enough food to get the needed nutrients. This is usually due to loss of appetite, poor absorption of nutrients, and changes in the way food is utilised in the body resulting from HIV and frequent infections.

2. The picture above

- The poor intake of food leads to loss of weight, body weakness, nutrient deficiencies and poor nutritional status.
- The poor nutritional status weakens the body's ability to fight diseases even further, and increases vulnerability to infections and to the impact of HIV.

3. The picture far right

- HIV destroys the body's natural ability to fight disease and infection.
- As a result the body's ability to fight infections is greatly reduced.




4. The picture below

- With a weakened ability to fight infections, the body becomes vulnerable to infections that normally may not have affected the person.
- The frequent infections make the body weaker, and lead to faster progression from HIV to AIDS.

Same person

She has lost weight
and looks sick.

She is healthier.



THEME 1

GOOD NUTRITION IS IMPORTANT FOR THE WELL-BEING OF PLWHA


1.3

GOOD NUTRITION DELAYS PROGRESSION OF HIV TO AIDS

- I. Ask the client to describe and explain the picture:
 - It is the same person. In one picture she has lost weight and looks sick. In the other picture she is healthy.
- II. Explain to the client that a PLWHA:
 - Can live a healthier life, free from frequent infections.
 - Can look well-nourished and be well-nourished.
- III. Explain that to live a healthier life and be well-nourished a PLWHA must take special care of his/her nutritional needs by:
 - Eating well,
 - Preventing infections,
 - Maintaining physical activity,
 - Managing diet related HIV/AIDS symptoms, such as diarrhoea.

Ask the client if he/she has any questions relating to any of the recommended practices. Use the following sets of cards to provide more information about the recommended practices.

It is important for PLWHA to eat a variety of food types to meet nutritional needs.



THEME 2 EATING WELL FOR PLWHA

2.1 EAT MEALS THAT HAVE A VARIETY OF FOOD TYPES

- I. Ask about the client's understanding of the need for food from each of the food types:
 - Energy giving foods (top and right of chart) - Provide the body with power, strength and energy to function. (e.g. cereals, tubers, oils)
 - Body building foods (bottom of chart) - Needed for building of muscles, cells, body defense system and enzymes. (e.g. animal products, nuts, beans)
 - Protective foods (left of chart) - Strengthen the body's ability to fight disease, help in cleansing body of toxins and by-products of body functions. (e.g. vegetables, fruits)
- II. Explain that it is important for a PLWHA to eat well to meet the body's nutritional needs for proper functioning, growth, and repair and for maintenance of health.
 - Different types of foods play different roles in the body. Therefore the body needs different types of foods to be able to meet its nutritional needs.
 - A PLWHA has increased nutritional needs resulting from the HIV infection. In some cases medication may also change nutritional needs.
 - Failing to meet the body's nutritional needs will lead to poor nutritional status. This will further weaken one's ability to fight diseases and will make one lose weight.
 - Eating well helps a PLWHA to meet their body's nutritional needs. It also can help medication work better.
- III. Explain that eating well means eating in adequate amounts of the different varieties of foods required by the body.
 - A good meal should include foods from the three food groups and a drink.
 - It is important to drink plenty of fluids, especially cool boiled water (at least 4 cups/8 glasses a day) preferably in between meals and after meals.
 - A PLWHA should have a good meal at least 3 times a day, plus snacks.
- IV. Discuss with the client food combinations that are available in his/her community. Ensure that they include the three food types plus clean water.

Ask the clients about questions she/he may have about what has been discussed.

Intake of essential nutrients can be increased by eating fortified foods.




THEME 2 EATING WELL FOR PLWHA

2.2 EAT FORTIFIED FOODS TO INCREASE INTAKE OF ESSENTIAL NUTRIENTS

- I. Explain to the client that intake of essential nutrients can be increased by eating fortified foods where available and affordable.
Fortified foods have added vitamins, minerals or other nutrients to improve their nutritional value.
- II. Ask the client to name fortified foods in the picture.
 - Salt fortified with iodine,
 - Commercially produced/packaged cereal flours like maize, millet, soya (fortified with the B-vitamins and zinc),
 - Milk products and yoghurt, mainly fortified with calcium and vitamin A,
 - Oils and fats like margarine and vegetable oil, mainly fortified with vitamins A and E,
 - Bottled fruit juices fortified with vitamin C (not in picture). Avoid quenchers since they simply contain coloured water, sugar and flavour.
- III. Ask the client if he/she knows more examples of fortified foods. Find out which foods the client can afford and is able to eat. Reinforce positive attitudes and practices. Clarify any myths the client may have about eating fortified foods.
- IV. Tell the client to always remember to read product labels (or if not possible, to ask) for indication of added nutrients and expiry dates.
- V. Tell the client that additional nutrient requirements can also be obtained by taking a multivitamin supplement.
- VI. Explain that multivitamin supplements should only be taken after consultation with a health worker.

STRESS: MULTIVITAMIN SUPPLEMENTS ARE ONLY ADDITIONS TO FOOD AND MUST NEVER REPLACE FOOD



PLWHA need to eat more in order to meet their increased nutritional needs.


2.3 INCREASE ENERGY INTAKE BY EATING MORE

- I. Explain to the client the importance of increasing intake of energy.
 - When one has HIV the body's requirement for energy is increased.
 - PLWHA need to eat more food than a person without HIV/AIDS. The healthiest way to meet the increased energy requirements is to eat more food from all food groups and maintain a balanced diet.
 - If the body's requirement for energy is not met, the body will break down fat and muscle tissue, leading to weight loss.

- II. Ask the client how they can increase intake of energy
 - Increase food intake by increasing the amount of food served.
 - Increase the frequency of meals and snacks.
 - Make every bite count by having foods that are rich in energy or by adding oil or sugar to food. However, large amounts of sugar and oil can be unhealthy as explained below.

- III. **IMPORTANT!** Caution the client to limit the following:
 - Amount of sugars, especially refined sugars. When taken in large amounts, sugars are stored as fat in the body. This can lead to diabetes, heart disease and liver damage. Sugar should be avoided if one has oral thrush.
 - Intake of fats and oils, especially from animal products. When taken in large amounts, they can cause high levels of fat in the blood, which can lead to heart disease, especially for PLWHA taking certain ARVs. Oil should be avoided if one has diarrhoea.
 - Intake of drinks like tea and coffee. These may affect absorption of nutrients and intake of other foods.

Meals and snacks
over a 24 hour
period



Discuss constraints to eating well and how to address them



THEME 2


EATING WELL FOR PLWHA

2.5 ADDRESS CONSTRAINTS TO EATING WELL

- Ask the client about problems and constraints they face in eating a variety of foods and increasing food intake as discussed.
- Explore and select options for addressing these problems. Below is a table with examples of problems that may be faced by a PLWHA and possible solutions.

If the client(s) reports	Then advise to:
Lack of time to prepare.	<ul style="list-style-type: none"> ▪ Use time saving methods such as katogo (mixing foods in one dish rather than cooking separate dishes) or mugoyo ▪ Double steaming or soaking dry legumes over night ▪ Buy a variety of already cooked foods (but not junk).
Lack of money to buy and include a variety foods in meals.	<ul style="list-style-type: none"> ▪ Backyard farming ▪ Make changes to allocation of household food budget to increase nutrition benefit ▪ Explore linking to community support groups
Crops recommended are normally not grown.	<ul style="list-style-type: none"> ▪ Grow these crops even on small scale in home garden
No one available to help buy, collect and prepare different foods.	<ul style="list-style-type: none"> ▪ Adopt time-saving cooking methods like mixing foods in one dish ▪ Refer to groups supporting food security
Traditional food taboos. How food is allocated within the home.	<ul style="list-style-type: none"> ▪ Encourage gradual change of attitude ▪ Involve family members in deciding on possible changes ▪ Seek alternatives
Seasonal variation in availability of food.	<ul style="list-style-type: none"> ▪ Use watering and mulching. ▪ Store foods during bumper harvests ▪ Refer to groups supporting food security
Loneliness during meals.	<ul style="list-style-type: none"> ▪ Consider involving caretakers to provide support and company for and during meal times. ▪ Refer for psycho-social counselling.
HIV related symptoms such as anorexia and oral thrush.	<ul style="list-style-type: none"> ▪ Advise on diet related management of symptoms using cards in Theme 4. ▪ Simple home medical management ▪ Refer for medical care if serious.

Clean home environment



Food hygiene and safety is one of the main ways of preventing infection



THEME 3 PREVENTING INFECTIONS


3.2 PRACTICE FOOD HYGIENE AND SAFETY

- I. Explain that food hygiene and safety is one of the main ways of preventing infection in PLWHA:
 - Food can be a source of infection if it is not properly handled, prepared and kept.
 - Most infections that arise out of poor food hygiene and safety practices are associated with diarrhoea and vomiting. These reduce intake and absorption of food, thus compromising nutritional status.

- II. Discuss vital food hygiene and safety practices in the picture:
 - Wash hands thoroughly before handling, preparing and eating foods.
 - Keep food and drinking water covered and stored away from insects, flies, rats and other animals.
 - Wash fruits and vegetables with clean water before eating, cooking or serving.
 - Use clean, safe water for food preparation.
 - Wash the food preparation area and eating and cooking utensils, and keep them clean.
 - Keep cooked food away from contact with raw food.

- III. Emphasise that the following should be avoided:
 - Eating mouldy, spoilt or rotten food.
 - Raw eggs or foods that contain raw eggs.
 - Raw fish.
 - Meat that is not well cooked.
 - Juices made with water that has not been previously boiled.

Regular check-ups are important for early identification of infections




THEME 3 PREVENTING INFECTIONS

3.3 HAVE REGULAR CHECK-UPS

- I. Explain that there are other actions a PLWHA can take to avoid getting infections. These include:
 - Practicing abstinence or safe sex.
 - Complying with recommended immunization and medical treatment for PLWHA.
 - Having regular visits to a health/HIV/AIDS care and support centre.
- II. Explain that it is necessary for PLWHA to periodically visit a health/HIV/AIDS care and support centre. PLWHA should visit such facilities for:
 - Periodic medical check-ups for any infections to ensure early management and treatment.
 - Weight monitoring to ensure right action is taken early in case one is underweight or overweight.
 - Services that may be necessary (e.g. immunizations, deworming, ART, treatment of other infections, and supplementation).
 - Information or skills that may be useful in managing HIV.
- III. Explain that in addition, PLWHA should seek medical care for diseases they may experience.
 - Specific health problems may include opportunistic infections, but also other illnesses like malaria.
 - Presence of all new and old symptoms should be explained to a health care provider on every visit to a care giving facility.
- IV. Discuss and allay any fears the PLWHA may have about visiting health/HIV/AIDS care facilities.

A woman takes a walk to maintain physical activity



Have a warm drink soon after waking in the morning or shortly before going to bed to help stimulate appetite.



Talk to friends to
help handle
depression



Oral thrush: A woman cleaning her mouth with cotton using salty water



5.3 MOUTH SORES AND ORAL THRUSH

- I. Explain that mouth sores or thrush are a common opportunistic infection in PLWHA that can affect eating.
- II. Discuss with the client the importance of cleaning the mouth to manage painful sores in the mouth:
 - Cleaning helps to prevent infection, helps to stop the infection from spreading, and promotes the healing process.
 - To clean, use cotton wool with mildly salty warm water. If possible, rinse the mouth with 1 teaspoon baking soda mixed in a glass of warm boiled water.
 - Clean at least twice a day: morning and evening, and preferably after every meal.
- III. Describe the dietary measures that can be used to relieve sores in the mouth.
 - Eat fermented products, e.g. yoghurt.
 - Eat soft foods such as mashed food, soups and juices.
 - Drink liquids using a straw to ease painful swallowing.
 - Eat ripe pawpaw to help heal the wounds.
 - Drink or rinse mouth with sour water. (To make sour water, soak grains/cereals for 2-3 days: 1 cup grain, and 3 cups of cool boiled water. Cover while fermenting).
 - Avoid acidic foods like lemon and oranges.

A man taking his
drugs with a snack



A woman picking dark green leafy vegetables in a makeshift garden



THEME 5 MANAGE DIET RELATED HIV/AIDS SYMPTOMS

5.5 ANAEMIA

- I. Tell the client that anaemia is common in PLWHA and may contribute to general body weakness. It weakens the body's ability to fight infections.
- II. Explain that eating foods that are rich in iron can help prevent and treat anemia.
- III. Identify and discuss with the client foods rich in iron that are accessible to the client.
 - Dark green leafy vegetables (spinach, sukuma wiki, dodo, nnakati, etc.)
 - Animal foods such as eggs, fish, and meat.
 - Avocado leaves, purple hibiscus leaves, cassava leaves, potato leaves.
 - Eating fruits rich in vitamin C like oranges and mangoes is important because vitamin C helps the body to absorb iron.
- IV. Emphasise to the client that he/she needs to:
 - Have haemoglobin checked at least every three months.
 - Be de-wormed periodically, at least every six months.
 - Avoid drinking coffee/tea with meals and avoid drinking too much coffee/tea because it reduces the absorption of iron by the body.
 - Treat malaria as soon as noticed.
 - Use iron supplements BUT only after consulting and with advice from a health provider.
- V. Refer the client to a nutritionist or a health worker if the cause of anaemia is not likely to be diet related. (For example, the ARV zidovudine can cause anaemia.)

Managing
diarrhoea: A man
taking an ORS
drink

